

The Role of DLG5 in Alzheimers Disease

<https://www.neurodegenerationresearch.eu/survey/the-role-of-dlg5-in-alzheimers-disease/>

Principal Investigators

XU, HUAXI

Institution

SANFORD BURNHAM PREBYS MEDICAL DISCOVERY INSTITUTE

Contact information of lead PI

Country

USA

Title of project or programme

The Role of DLG5 in Alzheimers Disease

Source of funding information

NIH (NIA)

Total sum awarded (Euro)

491972.4771

Start date of award

15/08/2015

Total duration of award in years

2

Keywords

Acquired Cognitive Impairment... Aging... Alzheimer's Disease... Alzheimer's Disease including Alzheimer's Disease Related Dementias (AD/ADRD)... Behavioral and Social Science... Brain Disorders... Dementia... Neurodegenerative... Neurosciences... Translational Research

Research Abstract

? DESCRIPTION (provided by applicant): Alzheimer's disease (AD) is the most common form of dementia in the elderly featuring progressive memory loss and cognitive decline. Compelling evidence suggests that soluble A β oligomers are responsible for early synaptic disruption and cognitive decline in AD. In contrast to A β -derived synaptic loss, the takusan domain family member a1-takusan enhances synaptic integrity and synaptic activity through interactions with PSD95 as demonstrated in our previous publication in Neuron. In our recent publication in the

Journal of Neuroscience, we have shown that overexpression of a1-takusan can alleviate A β -induced synaptic loss including decreased synaptic PSD-95 clustering, loss of dendritic spines, and depressed synaptic activity mediated by AMPA-type glutamate receptors. Interestingly, D2 is the most conserved sequence among all takusan family members, including human DLG5. For suitability purposes in downstream clinical applications (such as intranasal delivery) and simplified protein synthesis/purification, we generated a shorter DLG5-derived fragment hD2b comprising the human DLG5 D2 sequence and C-terminal PDZ-binding motif. Like a1-takusan, DLG5/hD2b alleviated A β -induced synaptic loss in cultured neurons as demonstrated in our previous publication. In preliminary studies, we have found that DLG5/hD2b overexpression reduced A β levels in conditioned medium from cultured N2a/APP^{swe} cells expressing human APP harboring amyloidogenic Swedish mutations. We further found that DLG5/hD2b interacted with β SGT, a co-chaperone protein and a negative regulator of HSP70, raising the possibility that DLG5/hD2b overexpression reduced A β levels and alleviated A β -induced synaptic loss at least in part through its ability to enhance HSP70-mediated A β -degradation. mRNA levels of the A β -degrading component insulin degrading enzyme (IDE), was consistently elevated by DLG5/hD2b overexpression. These results suggest an alternative mechanism underlying DLG5-mediated synaptic protection in addition to previously identified PSD-95-mediated synaptic protection. This hypothesis will be carefully tested in this project. In specific aim 1, we will determine the correlation between the level of DLG5 reduction and severity of dementia in AD patients and investigate the molecular mechanism underlying DLG5-mediated A β -reduction and synaptic protection. The role of β SGT/HSP70 chaperone proteins and A β -degrading enzymes in these actions will also be examined. In specific aim 2, we will determine the in vivo role of DLG5 as well as DLG5-mediated SGT/HSP70 chaperone activity in reducing amyloid pathology and cognitive deficits by performing immunoblotting, histochemistry, electrophysiology, and behavior tests in Tg2576 AD mice. If successful, our work will provide valuable information on the role of DLG5/hD2b against AD pathogenesis and establish groundwork for future development of DLG5-based therapeutics in AD.

Further information available at:

Types:

Investments < €500k

Member States:

United States of America

Diseases:

N/A

Years:

2016

Database Categories:

N/A

Database Tags:

N/A